

# WHAT TO PLANT WHEN

FOR THE ADELAIDE PLAINS (WARM TEMPERATURE CLIMATE)

Months to sow seeds	January	February	March	April	May	June	July	August	September	October	November	December
BROAD BEANS				█	█	█	█					
BASIL	█	█							█	█	█	█
BEANS, bush	█	█	█	█	█	█	█	█	█	█	█	█
BEANS, climbing	█								█	█	█	█
BETROOT	█	█	█	█	█	█	█	█	█	█	█	█
BROCCOLI		█	█	█	█	█	█	█	█	█	█	█
BRUSSELS SPROUTS*	█	█	█	█	█	█	█	█	█	█	█	█
CABBAGE	█	█	█	█	█	█	█	█	█	█	█	█
CAPSICUM								█	█	█	█	█
CARROT	█	█	█	█	█	█	█	█	█	█	█	█
CAULIFLOWER	█	█	█	█	█	█	█	█	█	█	█	█
CELERIAC								█	█	█	█	█
CELERY			█	█	█	█	█	█	█	█	█	█
CHEVRIL	█	█	█	█	█	█	█	█	█	█	█	█
CHILLI								█	█	█	█	█
CHIVES	█	█	█	█	█	█	█	█	█	█	█	█
CORIANDER			█	█	█	█	█	█	█	█	█	█
CORN	█									█	█	█
CUCUMBERS	█	█	█	█	█	█	█	█	█	█	█	█
DILL			█	█	█	█	█	█	█	█	█	█
EGGPLANT								█	█	█	█	█
ENDIVE			█	█	█	█	█	█	█	█	█	█
FENNEL			█	█	█	█	█	█	█	█	█	█
GARLIC			█	█	█	█	█	█	█	█	█	█
HARD SQUASH								█	█	█	█	█
KALE	█	█	█	█	█	█	█	█	█	█	█	█
KOHLRABI		█	█	█	█	█	█	█	█	█	█	█
LEEKS	█	█	█	█	█	█	█	█	█	█	█	█
LETTUCE	█	█	█	█	█	█	█	█	█	█	█	█
MACHE			█	█	█	█	█	█	█	█	█	█
MELONS								█	█	█	█	█
MINT	█	█	█	█	█	█	█	█	█	█	█	█
MIZUNA	█	█	█	█	█	█	█	█	█	█	█	█
MUSTARD RED	█	█	█	█	█	█	█	█	█	█	█	█
ONIONS, storage			█	█	█	█	█	█	█	█	█	█
PAK CHOI			█	█	█	█	█	█	█	█	█	█
PARSLEY								█	█	█	█	█
PARSNIP	█	█	█	█	█	█	█	█	█	█	█	█
PEAS				█	█	█	█	█	█	█	█	█
POTATOES								█	█	█	█	█
PUMPKINS									█	█	█	█
RADICCIO			█	█	█	█	█	█	█	█	█	█
RADISH	█	█	█	█	█	█	█	█	█	█	█	█
ROCKET	█	█	█	█	█	█	█	█	█	█	█	█
SALAD ONIONS	█	█	█	█	█	█	█	█	█	█	█	█
SILVERBEET	█	█	█	█	█	█	█	█	█	█	█	█
SPINACH		█	█	█	█	█	█	█	█	█	█	█
SPRING ONIONS	█	█	█	█	█	█	█	█	█	█	█	█
SWEDE			█	█	█	█	█	█	█	█	█	█
TATSOI	█	█	█	█	█	█	█	█	█	█	█	█
TOMATO	█	█	█	█	█	█	█	█	█	█	█	█
TURNIPS	█	█	█	█	█	█	█	█	█	█	█	█
ZUCCHINI								█	█	█	█	█

## Number of sowings and frequency

- █ Once (\*better suited to hills)
- █ Twice - Autumn and Spring
- █ Twice - once each season
- █ 1 to 2 - 2 months apart
- █ 2 to 3 - 3 weeks apart
- █ 2 to 3 - 4 weeks apart
- █ 2 to 3 - 6 weeks apart
- █ 2 to 3 - 8 weeks apart
- █ 2 to 3 - 2 months apart
- █ Multiple - 2 weeks apart
- █ Multiple - 2-3 weeks apart
- █ Multiple - 4 weeks apart
- █ Multiple - 8 weeks apart
- █ Multiple - every month
- █ Multiple - every 6 weeks

### Notes:

Plant families like the Solanaceae (tomato, capsicum, eggplant etc.) and the Cucurbitaceae (pumpkin, zucchini, squash, cucumber etc.) may need frost protection early in the season (Aug - Sept).

### Sources:

- Diggers - "The Australian Fruit and Vegetable Garden, Zone 10" (warm coastal)
- Peter Bennett - "Organic Gardening" (2006 - 7th ed)
- Norman de Vaus - "Better Vegetable Growing" (1973), based on South Australia Department of Agriculture table for Adelaide Plains
- Elliot Coleman "Four Season Harvest" (1999)
- Allsun Farm "Growing Annual Vegetables" CD-ROM

Compiled by Nat Wiseman, Village Greens of Willunga Creek.

Brought to you by:

**GREEN ADELAIDE**

